

AD \_\_\_\_\_

ARMY PROJECT ORDER NO: 91PP1805

TITLE: LONGTERM FOLLOW-UP OF PATIENTS IN CSP #298 "TREATMENT OF PATIENTS WITH 'ACQUIRED IMMUNE DEFICIENCY SYNDROME (AIDS) AND AIDS RELATED COMPLEX'"

PRINCIPAL INVESTIGATOR: Michael S. Simberkoff, M.D.

CONTRACTING ORGANIZATION: Veterans Administration Medical Center  
Cooperative Studies Program (121A4)  
819 Vermont Avenue NW  
Washington, DC 20420

REPORT DATE: October 1, 1994

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick  
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19941201 039

DTIC QUALITY INSPECTED 5

# REPORT DOCUMENTATION PAGE

FORM 298-102

OMB No. 0704-0186

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0186), Washington, DC 20503.

|  |   |  |   |   |  |
|--|---|--|---|---|--|
| 1. AGENCY USE ONLY (Leave blank)   |   | 2. REPORT DATE<br>October 1, 1994                          |   | 3. REPORT TYPE AND DATES COVERED<br>Final Report (4/1/91 - 4/30/94) |  |
| 4. TITLE AND SUBTITLE<br>Longterm Follow-up of Patients in CSP #298 "Treatment of Patients with 'Acquired immune Deficiency Syndrome (AIDS) and AIDS Related Complex'"   |   |  |   | 5. FUNDING NUMBERS<br>Army Project Order<br>91PP1805                |  |
| 6. AUTHOR(S)<br>Michael S. Simberkoff, M.D.<br>Pamela M. Hartigan, Ph.D.<br>John D. Hamilton, M.D.   |   |  |   |   |  |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>Veterans Administration Medical Center<br>Cooperative Studies Program (121A4)<br>819 Vermont Avenue NW<br>Washington, DC 20420   |   |  |   | 8. PERFORMING ORGANIZATION<br>REPORT NUMBER                         |  |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br>U.S. Army Medical Research and Materiel Command<br>Fort Detrick<br>Frederick, MD 21702-5012   |   |  |   | 10. SPONSORING/MONITORING<br>AGENCY REPORT NUMBER                   |  |
| 11. SUPPLEMENTARY NOTES  |   |  |   |   |  |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT<br>Approved for public release; distribution unlimited  |   |  |   | 12b. DISTRIBUTION CODE  |  |
| 13. ABSTRACT (Maximum 200 words)<br><br>To evaluate the benefits of early (E) vs. Later (L) ZDV therapy, we enrolled 338 symptomatic HIV+ patients with CD4 counts of 200-500/cmm into a randomized, double-blind trial. Patients were assigned to initial ZDV 1,500 mg/day (E: 168 pts) or initial placebo (L: 170 pts). The latter were switched to ZDV after progression to AIDS or to a CD4 count <200/cmm. All pts were offered ZDV 500 mg/day following completion of blinded follow-up. Blinded and final follow-up were completed in January 1991 and 1994, respectively. Median follow-up was 52 months. Progressions to AIDS occurred in 67 and 84 pts in the E and L therapy groups, respectively (rel. risk, 0.73; 95% CI, 0.52-1.01; p = 0.054). Death occurred in 74 and 73 pts in the E and L therapy groups, respectively (rel. risk, 0.99; 95% CI, 0.72-1.37; ns). We conclude that early compared with later ZDV provides marginal protection against progression to AIDS but no survival benefit. |   |  |   |   |  |
| 14. SUBJECT TERMS<br>AIDS, HIV Zidovudine, treatment   |   |  |   | 15. NUMBER OF PAGES   |  |
|  |   |  |   | 16. PRICE CODE  |  |
| 17. SECURITY CLASSIFICATION<br>OF REPORT<br>Unclassified   | 18. SECURITY CLASSIFICATION<br>OF THIS PAGE<br>Unclassified | 19. SECURITY CLASSIFICATION<br>OF ABSTRACT<br>Unclassified | 20. LIMITATION OF ABSTRACT<br>Unlimited |   |  |

## TABLE OF CONTENTS

|       |   |    |
|-------|---|----|
| I.    | Introduction  | 2  |
| II.   | Methods   | 2  |
| III.  | Results   | 5  |
| IV.   | Conclusions   | 6  |
| V.    | Addendum (specimens sent to Walter Reed)                  | 8  |
| VI.   | References  | 8  |
| VII.  | Bibliography of work published and abstracts presented    | 10 |
| VIII. | List of personnel receiving pay from the contract support | 13 |
|       | Tables 1-4  | 15 |
|       | Figures 1-3   | 19 |

|                    |                                     |
|--------------------|-------------------------------------|
| Accession For      |                                     |
| NTIS CRA&I         | <input checked="" type="checkbox"/> |
| DTIC TAB           | <input type="checkbox"/>            |
| Unannounced        | <input type="checkbox"/>            |
| Justification      |                                     |
| By                 |                                     |
| Distribution /     |                                     |
| Availability Codes |                                     |
| Dist               | Avail and/or<br>Special             |
| A-1                |                                     |

## I. INTRODUCTION

The Agency for Health Care Policy and Research, a division of the U.S. Public Health Service, recommends zidovudine as the first anti-retroviral treatment for patients who have CD4+ counts less than 500/mm<sup>3</sup>, with or without symptoms (1). This recommendation is based on randomized, placebo-controlled studies that showed zidovudine to be effective in preventing progression to AIDS in this group of patients (2-4) and on retrospective cohort analyses that showed longer survival in patients who received zidovudine early in their course of illness (5,6). Nevertheless, there are conflicting data. Concorde, the largest trial published to date comparing early to delayed zidovudine therapy, showed no benefit to early therapy (7).

In 1987, the Cooperative Studies Program (CSP) of the Department of Veterans Affairs initiated a randomized trial comparing early with later zidovudine therapy in symptomatic HIV-infected patients. Double-blinded follow-up of the original cohort of 338 patients was terminated in January 1991, and the results of that portion of the study were published (8). At completion of the double-blinded follow-up, study subjects were offered continued "unblinded" zidovudine treatment in a dose of 500 mg/day at participating medical centers, and they were followed for an additional 3 years until January 1994. The results of the combined follow-up of the cohorts originally randomized to early or later zidovudine are detailed in this report.

## II. METHODS

*Patient population.* This report combines the results of Part I and Part II of the

study; Part I (the blinded position) was from January 1987 to January 1991, and Part II (the unblinded position) was from January 1991 to January 1994. The criteria for selection of the original group of 338 patients were described previously (8).

The protocol and consent forms for both Part I and Part II of this study were approved by the Human Rights Committee of the Cooperative Studies Program and by the Investigational Review Board at each participating hospital. All patients gave written, informed consent.

*Treatment regimens.* The treatment regimens were as follows:

Early zidovudine: zidovudine treatment beginning at randomization in a dose of 1,500 mg/day in divided doses;

Later zidovudine: initial placebo changed to zidovudine when the CD4+ count fell to  $<200/\text{mm}^3$  or an AIDS-defining event occurred.

Randomization of patients was completed in January 1990, and blinded follow-up was completed in January 1991. At the completion of blinded follow-up, all patients were offered zidovudine in a dose of 500 mg/day in divided doses. Thus, patients randomly assigned to later zidovudine treatment received placebo until they reached a clinical or immunological end point, defined below, or for a minimum period of 1 year.

*Criteria for response.* Death and AIDS were the clinical end points of this study. We used the 1987 Centers for Disease Control and Prevention (CDC) surveillance criteria for the diagnosis of AIDS throughout this study (9). Our definition of dementia was described previously (8). *AIDS-related death* was defined as death associated with a

current AIDS-defining condition. *Death with HIV progression* was defined as death preceded by AIDS or an HIV-associated illness (e.g., pneumococcal bacteremia and meningitis), increased symptoms, or a fall in CD4+ concentration to <200 cells per cubic millimeter. The immunological end point of this study was the fall in CD4+ concentration to less than 200 cells per cubic millimeter on two successive measurements done 6 or more weeks apart.

*Evaluation and follow-up.* Follow-up of patients was at least monthly during Part I and at least bimonthly during Part II of this study. A detailed history, physical examination, and laboratory studies were obtained at each of these evaluations.

*Data management and statistical analyses.* Data collected at participating sites were sent to the study co-chairman's office and the CSP Coordinating Center for review. Methods of data verification and entry were described previously (8). Comparability of treatment groups was assessed by a chi-square test or Fisher's exact test for discrete variables, or by Student's *t* test or Wilcoxon rank-sum test for continuous variables (10). Time to clinical and immunologic end points was estimated with Kaplan-Meier and proportional-hazards regression methods. Stratified log-rank tests and proportional hazards models were used to compare treatment groups and to estimate relative risks and confidence intervals (CIs), adjusting for the stratification by CD4+ cell concentration (11). Qualitative interactions were tested with the method developed by Gail and Simon (12). Throughout, we have reported the relative risks comparing early with later zidovudine treatment.

### III. RESULTS

*Patient characteristics.* The baseline characteristics of our original cohort of 338 patients were described in our earlier manuscript (8). Forty-four (44) patients died, and 15 patients were lost to follow-up during Part I of this study. There were 9 other patients from whom no data were collected during Part II. Thus, Part II had 270 patients, equally divided between the two treatment groups, from whom data were collected. Table 1 summarizes the baseline characteristics of the patients followed in Part II of the study. There were no differences between the treatment groups, nor were there significant differences between this group and the group that died or was lost during Part I of the study.

*Progression to AIDS.* CDC-defined AIDS occurred in 67 and 84 patients in the early- and later-therapy groups, respectively (Table 2). There were no significant differences in the occurrences of individual AIDS-defining diagnoses between the treatment groups. The number of patients with first AIDS events in each treatment group at yearly intervals of follow-up is shown in Table 3, Part A. The relative risk of progression to AIDS and the 95% CIs for early compared with later treatment are also shown. The relative risk for progression to AIDS ranged from 0.68 to 0.73. The 95% CIs just barely overlapped 1 at 3, 4, and 5 years of follow-up and at study conclusion. A Kaplan-Meier lifetable plot of the time to AIDS in the study population is shown in Figure 1. The median time to AIDS in the later-therapy group was 57.4 months whereas it was 68.2 months in the early-therapy group. This is a difference of almost 11 months. By stratified log-rank test, the  $p$  was 0.054.

*Deaths.* There were 74 deaths in the early-therapy group and 73 deaths in the later-therapy group (Table 3, Part B). The relative risk of death for the early-therapy group was close to 1 throughout the study (Figure 2A). A Kaplan-Meier lifetable analysis of the time to death showed virtually identical curves in the two treatment groups. The median times to death was 65.8 months and 68.7 months in the early- and later-therapy groups, respectively. By stratified log-rank test, the  $p$  was 0.96 (Table 4). Of the 74 patients who died in the early-therapy group, 48 (65%) had AIDS, an additional 16 (22%) had an AIDS-related disease, and 10 died of AIDS-unrelated or unknown causes. Of the 73 patients who died in the later-therapy group, 64 (87.6%) had AIDS, an additional 3 (4%) had an AIDS-related disease, and 6 (8%) died of AIDS-unrelated or unknown causes (Figure 2B). For survival after an AIDS diagnosis, there was little difference between the two therapy groups. The median survival after AIDS in the early-therapy group was 12.9 months; it was 15.9 months in the later-therapy group ( $p = 0.64$ ).

*Immunologic progressions.* Absolute CD4+ counts increased in the early-therapy group (Figure 3). In this group, the mean absolute CD4+ remained above baseline for the first year of follow-up. In the later-therapy group, the CD4+ counts declined steadily. Statistically significant differences between the treatment groups persisted for over 3 years of follow-up.

#### IV. CONCLUSIONS

Our data indicate that early zidovudine therapy for patients with symptomatic



HIV infection and initial CD4+ counts between 200 and 500/mm<sup>3</sup> results in a significant, sustained CD4+ count elevation and provides marginal protection against progression to AIDS but no survival benefit. These findings are similar to those that we previously reported (8). However, by extending the period of follow-up for our patients, we have increased the number of events observed and narrowed the confidence intervals of our relative risk estimates.

Our findings that early zidovudine therapy reduced the risk of AIDS-defining events is similar to that reported in other prospective zidovudine-versus-placebo-controlled trials, including those reported by the AIDS Clinical Trials Group (3,4) and by the European-Australian Collaborative Group (13). It must be recognized, however, that many of the opportunistic infections observed in our study and in the others reported can be prevented by other forms of prophylaxis that are now widely used (1).

Our findings on death are similar to those reported by the Concorde Coordinating Committee (7). In the Concorde trial, as in ours, there was a trend toward an increased death rate in the early-therapy group. These findings tend to support the now more widely held view that prolonged anti-retroviral monotherapy may not be of benefit and, in some instances, may be detrimental to HIV-infected patients (14).

Despite optimistic statements, the optimum time for initiating anti-retroviral therapy remains a legitimate subject for debate among investigators (15), as well as a subject for discussion between patients and their physicians. Although there is hope that cycles or combinations of anti-retrovirals will reduce morbidity and prolong survival of HIV-infected patients, the clinical trials necessary to prove these clinical benefits have

not been completed.

## V. ADDENDUM (specimens sent to Walter Reed)

Another goal of the project was to collect and send to the Army research laboratory under the direction of Dr. Douglas Mayers an aliquot of lymphocytes from each blood specimen received at the study's reference laboratory at Duke University. In addition, an aliquot of the supernatant from all positive cultures was to be forwarded to the same laboratory. During the period of the study, these specimens were regularly assembled and sent as prescribed. The final shipment is currently being assembled and will be forwarded by October 15, 1994.

## VI. REFERENCES

1. Clinical Practice Guidelines, Number 7: Evaluation and management of early HIV infection. U.S. Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research. AHCPR Publication No. 94-0572.
2. Fischl MA, Richman DD, Grieco MH, et al: The efficacy of azidothymidine (AZT) in the treatment of patients with AIDS and AIDS-related complex: A double-blind, placebo-controlled trial. N Engl J Med 317:185-191, 1987.
3. Fischl MA, Richman DD, Hansen N, et al: The safety and efficacy of zidovudine (AZT) in the treatment of mildly symptomatic human immunodeficiency virus

- type 1 (HIV) infection: A double-blind, placebo-controlled trial. *Ann Intern Med* 112:727-737, 1990.
4. Volberding PA, Lagakos SW, Koch MA, et al: Zidovudine in asymptomatic human immunodeficiency virus infection: A controlled trial in persons with fewer than 500 CD4-positive cells per cubic millimeter. *N Engl J Med* 322:941-949, 1990.
  5. Moore RD, Hidalgo J, Sugland BW, Chaisson RE. Zidovudine and the natural history of the acquired immunodeficiency syndrome. *N Engl J Med* 1991; 324:1412-6.
  6. Graham NM, Zeger SL, Park LP, et al. The effects on survival of early treatment of human immunodeficiency virus infection. *N Engl J Med* 1992; 326:1037-42.
  7. Concorde Coordinating Committee. Concorde MRC/ANRS randomized double-blind controlled trial of immediate and deferred zidovudine in symptom-free HIV infection. *Lancet* 1994; 343:871-81.
  8. Hamilton JD, Hartigan PM, Simberkoff MS, et al. A controlled trial of early versus later zidovudine in symptomatic human immunodeficiency virus infection: Results of a Veterans Affairs Cooperative Study. *N Engl J Med* 1992; 326:437-43.
  9. Centers for Disease Control. Revision of the CDC surveillance case definition for acquired immunodeficiency syndrome. *MMWR* 1987;36:Suppl 1S:1-16.
  10. Peto R, Pike MC, Armitage P, et al. Design and analysis of randomized clinical trials requiring prolonged observation of each patient: II. Analysis and examples. *Br J Cancer* 1977;35:1-39.

11. Kalbfleisch JD, Prentice RG. The statistical analysis of failure time data. New York: John Wiley & Sons, 1980.
12. Gail M, Simon R. Testing for qualitative interactions between treatment effects and patient subsets. *Biometrics* 1985;41:361-72.
13. Cooper DA, Gatell JM, Kroon S, et al. Zidovudine in persons with asymptomatic HIV infection and CD4+ cell counts greater than 400 per cubic millimeter. *N Engl J Med* 1993; 329:297-303.
14. Caliendo AM, Hirsch MS. Combination therapy for infection due to human immunodeficiency virus, type 1. *Clin Infect Dis* 1994; 18:516-24.
15. Friedland GH. Early treatment for HIV: The time has come. *N Engl J Med* 1990;322:1000-2.

## VII. BIBLIOGRAPHY

### A. *Published articles*

1. Hamilton JD, Hartigan PM, Simberkoff MS, et al. A controlled trial of early versus late treatment with zidovudine in symptomatic human immunodeficiency virus infection: Results of the Veterans Affairs Cooperative Study. *N Engl J Med* 326:437-443, 1992.
2. Hamilton J. Late treatment of HIV infection: Is it better? *Infectious Agents and Disease* 1:156-162, 1992.
3. Economou TM, Hartigan PM. Linear time lines: A visual summary of critical patient data. *SAS Users Group International 17 Conference Proceedings*, pp. 1036-

- 1041, 1992.
4. Simberkoff M, et al. Letter to the editor: Effects of zidovudine therapy in minority and other subpopulation with early HIV infection. *JAMA* 267:2472-2473, 1992.
  5. Letter to the editor: Clinical trials: To terminate or not to terminate? *Ann Intern Med* April 15, 1992.
  6. Response to letter to editor: Early vs late zidovudine in HIV infected individuals. *N Engl J Med* 327:814-815, 1992.
  7. St. Clair MH, Hartigan PM, Andrews JC, et al. Zidovudine resistance, syncytium-inducing phenotype, and HIV disease progression in a case-control study. *J AIDS* 6:891-987, 1993. [To be abstracted in the *Year Book of Infectious Diseases*, 1995, Chicago, IL: Mosby Year Book, Inc. Also, article won the 1994 Howard Temin Award for the outstanding paper in clinical medicine]
  8. Simberkoff MS, Hartigan PM, Hamilton JD, et al. Ethical dilemmas in continuing a zidovudine trial after early termination of similar trials. *Controlled Clin Trials* 14:6-18, 1993.
  9. Oddone EZ, Cowper PA, Hartigan PM, et al.: A cost effectiveness analysis of early ZDV therapy. *Br Med J* 307:1322-1325, 1993.
  10. Gordin F, Hartigan P, Klimas NG, et al. Delayed-type hypersensitivity (DTH) skin tests are an independent predictor of HIV disease progression. *J Infect Dis* 169:893-897, 1994.
  11. Hamilton JD. When to treat HIV-infected patients, with what, and why? *Clinical*

*Performance and Quality Health Care* 2:115-117, 1994.

*B. Abstracts presented*

1. Hamilton JD. Long-term studies of AZT use. Paper presented at the "State of the Art Symposium," VIII International Conference on AIDS/III STD World Congress, Amsterdam, The Netherlands, July 22, 1992.
2. Gordin F, Hartigan P, Simberkoff M, et al. Delayed-type hypersensitivity (DTH) reactions are an independent predictor of progression of human immunodeficiency virus. Presented at the 32nd ICAAC, Anaheim, California, October, 1992.
3. Oddone EZ, Cowper P, Matchar D, et al. Markov process cost-effectiveness of early versus later zidovudine (AZT) treatment of symptomatic HIV infection. Presented at the Society of General Internal Medicine, May, 1992.
4. St. Clair M, Hartigan PM, Andrews J, et al. Matched progressor-nonprogressor study of zidovudine resistance and disease progression. Presented at the VIII International Conference on AIDS, July, 1992.
5. Simberkoff MS, Hartigan PM, Hamilton JD, et al. Longer follow-up of VA trial comparing early versus late AZT for symptomatic HIV infection. Presented at the VIII International Conference on AIDS, July, 1992.
6. Hamilton JD. Implications of early versus later treatment of symptomatic HIV infection: Report of the VA Cooperative study. Paper presented at the Institut Pasteur, Paris, France, March 5, 1993.
7. Hamilton JD. Implications of early versus later treatment of symptomatic HIV

infection: Report of the VA Cooperative study. Paper presented at the Karolinska Institut, Stockholm, Sweden, March 8, 1993.

8. Hamilton JD. Clinical care and prevention insights from early intervention research. Presentation to be made at the AMA Early Intervention Conference, which precedes the IX International Conference on AIDS, Berlin, Germany, June 6, 1993.
9. Bailey EM, Haren N, SanFelipo M, et al. Factors of participation and decision to remain in long term early/late AZT treatment clinical trial. Presented at the Association of Nurses in AIDS Care Conference, New York, NY, April 19, 1993.
10. Simberkoff M, Hartigan P, Hamilton J, et al. Final report on longterm followup of symptomatic HIV+ patients randomized to early vs later zidovudine (ZDV) Rx. To be presented at ICAAC, Orlando, Florida, October, 1994.

#### VIII. LIST OF PERSONNEL RECEIVING PAY FROM THE CONTRACT SUPPORT

##### A. Patient treatment centers:

*Manhattan VAMC:* Noreen Haren, R.N.; Eileen Bailey, R.N.

*Miami VAMC:* Gishlaine Paperwallar, R.N.; John Roussel, M.S.

microbiology (laboratory technician)

*Houston VAMC:* Adonna Peacock, R.N.; Joanne Mitchell

*Los Angeles VAMC:* Carol Silbar, R.N.; Felicitas Lorenzo, R.N.

*San Francisco VAMC:* Roland Jalbert, R.N., Manon Maravich, R.N., Ann

Cotleur, B.S. (lab technician)

*Washington, DC, VAMC:* Patricia Ackerson, R.N.

*Durham, NC, Cochairs Office:* Patricia Spivey, Study Coordinator; Clorine

Chasten, Isabel Routh, Lucy Skinner, secretaries

*Duke Virology Laboratory, Durham, NC:* Debbie Wakefield (laboratory

technician), Janet Ottinger (laboratory technician),

Caroline Hinton (laboratory technician), Tom

Matthews, Ph.D. (laboratory supervisor)

*Baltimore VAMC Reference Laboratory:* Mark Rossen (laboratory

technician)



Table 1

## Baseline Characteristics of Patients Followed in Part II

|                                      | <u>Zidovudine Treatment</u> |               |
|--------------------------------------|-----------------------------|---------------|
|                                      | Early (N=135)               | Later (N=135) |
| Stratum - No. (%) of patients        |                             |               |
| 1                                    | 33 (24)                     | 34 (25)       |
| 2                                    | 102 (76)                    | 101 (75)      |
| Age at randomization (yr)            | 40.4                        | 40.2          |
| Sex - No. of patients                |                             |               |
| Male                                 | 133                         | 134           |
| Female                               | 2                           | 1             |
| Race - No. (%) of patients           |                             |               |
| Non-Hispanic white                   | 92 (68)                     | 94 (69)       |
| Black or Hispanic                    | 37 (27)                     | 39 (29)       |
| Risk group - No. (%) of patients     |                             |               |
| Homosexual or bisexual               | 87 (65)                     | 82 (61)       |
| Intravenous drug use                 | 20 (15)                     | 26 (19)       |
| Homosexual and IVDU                  | 14 (10)                     | 13 (10)       |
| Other                                | 14 (10)                     | 14 (10)       |
| No. of symptoms (Mean)               | 3.0                         | 3.3           |
| CD4+ count/mm <sup>3</sup> - Mean±SD | 353.4±92.7                  | 362±80.7      |
| Karnofsky score                      | 8.9                         | 8.9           |

Table 2  
Progression to AIDS - First Diagnoses

| Diagnosis                  | <u>Zidovudine Treatment</u> |      |
|----------------------------|-----------------------------|------|
|                            | Early                       | Late |
| Kaposi's sarcoma           | 8                           | 12   |
| PCP                        | 21                          | 25   |
| Esophageal candidiasis     | 16                          | 10   |
| Disseminated MAC infection | 4                           | 4    |
| Toxoplasmosis              | 2                           | 3    |
| Cryptococcosis             | 1                           | 1    |
| Cryptosporidiosis          | 0                           | 1    |
| PML                        | 1                           | 4    |
| Lymphoma                   | 2                           | 7    |
| CMV (retinitis or GI)      | 4                           | 7    |
| Disseminated herpes        | 1                           | 1    |
| Extra-pulmonary TB         | 1                           | 0    |
| Dementia                   | 2                           | 7    |
| Wasting                    | 4                           | 2    |
| Total                      | 67                          | 84   |

Table 3

## Number of Patients and Relative Risk of Progression to AIDS and Death

## A. Progression to AIDS

| Follow-up | <u>Zidovudine Treatment</u> |      | Relative risk (95% CI) | <i>p</i> |
|-----------|-----------------------------|------|------------------------|----------|
|           | Early                       | Late |                        |          |
| 1 Year    | 11                          | 15   | 0.69 (0.32 - 1.52)     |          |
| 2 Years   | 25                          | 35   | 0.68 (0.95 - 1.13)     |          |
| 3 Years   | 40                          | 55   | 0.69 (0.46 - 1.09)     |          |
| 4 Years   | 54                          | 71   | 0.71 (0.50 - 1.00)     |          |
| 5 Years   | 54                          | 78   | 0.73 (0.52 - 1.02)     |          |
| Final     | 67                          | 84   | 0.73 (0.52 - 1.01)     | 0.054    |

## B. Death

| Follow-up | <u>Zidovudine Treatment</u> |      | Relative risk (95% CI) | <i>p</i> |
|-----------|-----------------------------|------|------------------------|----------|
|           | Early                       | Late |                        |          |
| 1 Year    | 5                           | 3    | 1.62 (0.39 - 6.72)     |          |
| 2 Years   | 11                          | 8    | 1.35 (0.54 - 3.36)     |          |
| 3 Years   | 34                          | 22   | 1.51 (0.88 - 2.59)     |          |
| 4 Years   | 50                          | 51   | 0.97 (0.65 - 1.44)     |          |
| 5 Years   | 68                          | 63   | 1.06 (0.74 - 1.50)     |          |
| Final     | 74                          | 73   | 0.99 (0.72 - 1.37)     | 0.962    |

Table 4  
Relation Between AIDS, HIV-Related Disease, and Death

|                      | <u>Zidovudine Treatment</u> |            |
|----------------------|-----------------------------|------------|
|                      | Early                       | Later      |
| Prior AIDS           | 48 (64.9%)                  | 64 (87.7%) |
| HIV-related disease  | 16 (21.6%)                  | 3 (4.1%)   |
| Unrelated or unknown | 10 (13.5%)                  | 6 (8.2%)   |
| Total                | 74 (100%)                   | 73 (100%)  |

## LEGENDS

- Figure 1. Estimated Kaplan-Meier distribution of the time to AIDS, according to study group.
- Figure 2A. Estimated Kaplan-Meier distribution of the time to death, according to study group.
- Figure 2B. Estimated Kaplan-Meier distribution of the time from an AIDS diagnosis to death, according to study group.
- Figure 3. Mean ( $\pm$ SE) changes from baseline in absolute CD4+ counts, according to study group.




Figure #1




Figure 2A

# SURVIVAL AFTER AN AIDS DIAGNOSIS




Figure 2B



# CD4+ COUNT OVER TIME

Means  $\pm$  SE




Figure 3